

# Incompleteness Theorems for Observables in General Relativity

Aristotelis Panagiotopoulos

Kurt Gödel Research Center  
University of Vienna



Research supported by DMS-2154258

This is joint work with  
Marios Christodoulou (IQOQI)



George Sparling (UPitt)



# Table of Contents

1 Spacetimes

2 Observables

3 The Proof

4 Future Directions

# Lorentzian metrics: abstractly


## Lorentzian metrics: abstractly

Let  $M$  be smooth 4-dimensional manifold. A **Lorentzian metric** on  $M$  is “a symmetric and  $(1, 3)$ -signature **section**  $g$  of the bundle  $(TM \otimes TM)^* \rightarrow M$ ”

# Lorentzian metrics: abstractly

Let  $M$  be smooth 4-dimensional manifold. A **Lorentzian metric** on  $M$  is  
“a symmetric and  $(1, 3)$ -signature **section**  $g$  of the bundle  $(TM \otimes TM)^* \rightarrow M$ ”


$$g(\vec{V}, \vec{W})_p := \text{the “inner product” of } \vec{V}, \vec{W} \text{ at } p \in M$$



# Lorentzian metrics: abstractly

Let  $M$  be smooth 4-dimensional manifold. A **Lorentzian metric** on  $M$  is  
“a symmetric and  $(1, 3)$ -signature **section**  $g$  of the bundle  $(TM \otimes TM)^* \rightarrow M$ ”


$$g(\vec{V}, \vec{W})_p := \text{the “inner product” of } \vec{V}, \vec{W} \text{ at } p \in M$$



# Lorentzian metrics: abstractly

Let  $M$  be smooth 4-dimensional manifold. A **Lorentzian metric** on  $M$  is  
“a symmetric and  $(1, 3)$ -signature **section**  $g$  of the bundle  $(TM \otimes TM)^* \rightarrow M$ ”

$$g(\vec{V}, \vec{W})_p := \text{the “inner product” of } \vec{V}, \vec{W} \text{ at } p \in M$$






# Lorentzian metrics: abstractly

Let  $M$  be smooth 4-dimensional manifold. A **Lorentzian metric** on  $M$  is “a symmetric and  $(1, 3)$ -signature **section**  $g$  of the bundle  $(TM \otimes TM)^* \rightarrow M$ ”

$$g(\vec{V}, \vec{W})_p := \text{the “inner product” of } \vec{V}, \vec{W} \text{ at } p \in M$$



*R. Penrose, “The Road to Reality”*

## Lorentzian metrics: concretely

A **Lorentzian metric** on  $\mathbb{R}^4$  is given by a smooth map  $g_{\mu\nu} : \mathbb{R}^4 \rightarrow \mathbb{R}^{4 \times 4}$

$$(x^0, x^1, x^2, x^3) \mapsto \begin{bmatrix} g_{00} & g_{01} & g_{02} & g_{03} \\ g_{10} & g_{11} & g_{12} & g_{13} \\ g_{20} & g_{21} & g_{22} & g_{23} \\ g_{30} & g_{31} & g_{32} & g_{33} \end{bmatrix}$$

with  $g_{\mu\nu}$  being a symmetric,  $(-, +, +, +)$ -signature matrix. We have

$$g = g_{\mu\nu} dx^\mu dx^\nu$$

## Lorentzian metrics: concretely

A **Lorentzian metric** on  $\mathbb{R}^4$  is given by a smooth map  $g_{\mu\nu} : \mathbb{R}^4 \rightarrow \mathbb{R}^{4 \times 4}$

$$(x^0, x^1, x^2, x^3) \mapsto \begin{bmatrix} g_{00} & g_{01} & g_{02} & g_{03} \\ g_{10} & g_{11} & g_{12} & g_{13} \\ g_{20} & g_{21} & g_{22} & g_{23} \\ g_{30} & g_{31} & g_{32} & g_{33} \end{bmatrix}$$

with  $g_{\mu\nu}$  being a symmetric,  $(-, +, +, +)$ -signature matrix. We have

$$g = g_{\mu\nu} dx^\mu dx^\nu$$

### Example

$$\text{If } \eta_{\mu\nu} := \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}, \quad \text{then } \eta = -dt^2 + dx^2 + dy^2 + dz^2$$

# Einstein field equations

By a **spacetime** we mean a Lorentzian metric  $g_{\mu\nu}: \mathbb{R}^4 \rightarrow \mathbb{R}^{4 \times 4}$ , satisfying:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = \frac{8\pi G}{c^4}T_{\mu\nu}$$

for some “physically relevant” **Stress-Energy tensor**  $T_{\mu\nu}$ .

$$g_{\mu\nu} \rightsquigarrow R_{\mu\sigma\nu}^{\rho} \rightsquigarrow R_{\mu\nu} \rightsquigarrow R$$

Compare to Poisson's equation for Newton's law of gravity:

$$\nabla^2\varphi = 4\pi G\rho$$

## Example

$$g_{\mu\nu} := 1/(2\omega^2) \left[ - (dt + e^x dy)^2 + dx^2 + 1/2 e^{2x} dy^2 + dz^2 \right]$$

$T_{\mu\nu}$  = “rotating dust” + “negative cosmological constant”




Figure. *Nméti, Madarász, Andréka, Andai* (after Hawking, Ellis)

## Example

$$g_{\mu\nu} := 1/(2\omega^2) [ - (dt + e^x dy)^2 + dx^2 + 1/2 e^{2x} dy^2 + dz^2 ]$$

$T_{\mu\nu}$  = “rotating dust” + “negative cosmological constant”




Figure. *Nm*

king, Ellis)

“Is the universe rotating yet?” K. Gödel

# Table of Contents

1 Spacetimes

2 Observables

3 The Proof

4 Future Directions

**Question.** Do  $g_{\mu\nu}$  and  $\tilde{g}_{\rho\sigma}$  represent different “geometries”?

$$g_{\mu\nu} := \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\tilde{g}_{\rho\sigma} := \begin{bmatrix} -1 & -\cos(x_1) & 0 & 0 \\ -\cos(x_1) & 1 - \cos^2(x_1) & 2x_2 & 0 \\ 0 & 2x_2 & 4x_2^2 + 1 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}$$



**Question.** Do  $g_{\mu\nu}$  and  $\tilde{g}_{\rho\sigma}$  represent different “geometries”?

$$g_{\mu\nu} := \begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

$$\tilde{g}_{\rho\sigma} := \begin{bmatrix} -1 & -\cos(x_1) & 0 & 0 \\ -\cos(x_1) & 1 - \cos^2(x_1) & 2x_2 & 0 \\ 0 & 2x_2 & 4x_2^2 + 1 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}$$

We say that  $g_{\mu\nu}$  and  $\tilde{g}_{\rho\sigma}$  are **diffeomorphic** and write  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma}$  if there exists a smooth change of coordinates  $x_\eta = x_\eta(\tilde{x}^\xi)$  so that

$$\tilde{g}_{\rho\sigma}(\tilde{x}^\xi) = \frac{\partial x^\mu}{\partial \tilde{x}^\rho} \frac{\partial x^\nu}{\partial \tilde{x}^\sigma} g_{\mu\nu}(x^\eta) \quad \text{for all } \tilde{x}^\xi.$$

Same geometry, different coordinate system...

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \simeq_{\text{diff}} \begin{bmatrix} -1 & -\cos(\tilde{x}_1) & 0 & 0 \\ -\cos(\tilde{x}_1) & 1 - \cos^2(\tilde{x}_1) & 2\tilde{x}_2 & 0 \\ 0 & 2\tilde{x}_2 & 4\tilde{x}_2^2 + 1 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}$$

Same geometry, different coordinate system...

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \simeq_{\text{diff}} \begin{bmatrix} -1 & -\cos(\tilde{x}_1) & 0 & 0 \\ -\cos(\tilde{x}_1) & 1 - \cos^2(\tilde{x}_1) & 2\tilde{x}_2 & 0 \\ 0 & 2\tilde{x}_2 & 4\tilde{x}_2^2 + 1 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}$$

Consider the change of coordinates  $x_\eta = x_\eta(\tilde{x}^\xi)$  given by:

$$x_0 = \tilde{x}_0 + \sin(\tilde{x}_1)$$

$$x_1 = \tilde{x}_1 + \tilde{x}_2^2$$

$$x_2 = \tilde{x}_2 - \tilde{x}_3$$

$$x_3 = \tilde{x}_3$$

Same geometry, different coordinate system...

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \simeq_{\text{diff}} \begin{bmatrix} -1 & -\cos(\tilde{x}_1) & 0 & 0 \\ -\cos(\tilde{x}_1) & 1 - \cos^2(\tilde{x}_1) & 2\tilde{x}_2 & 0 \\ 0 & 2\tilde{x}_2 & 4\tilde{x}_2^2 + 1 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}$$

Consider the change of coordinates  $x_\eta = x_\eta(\tilde{x}^\xi)$  given by:

$$\begin{array}{ll} x_0 = \tilde{x}_0 + \sin(\tilde{x}_1) & dx_0 = d\tilde{x}_0 + \cos(\tilde{x}_1)d\tilde{x}_1 \\ x_1 = \tilde{x}_1 + \tilde{x}_2^2 & dx_1 = d\tilde{x}_1 + 2\tilde{x}_2d\tilde{x}_2 \\ x_2 = \tilde{x}_2 - \tilde{x}_3 & dx_2 = d\tilde{x}_2 - d\tilde{x}_3 \\ x_3 = \tilde{x}_3 & dx_3 = d\tilde{x}_3 \end{array} \rightsquigarrow$$

$$\rightsquigarrow \begin{array}{ll} (dx_0)^2 = & (d\tilde{x}_0)^2 + 2\cos(\tilde{x}_1)d\tilde{x}_0d\tilde{x}_1 + \cos^2(\tilde{x}_1)(d\tilde{x}_1)^2 \\ (dx_1)^2 = & (d\tilde{x}_1)^2 + 4\tilde{x}_2d\tilde{x}_1d\tilde{x}_2 + 4\tilde{x}_2^2(d\tilde{x}_2)^2 \\ (dx_2)^2 = & (d\tilde{x}_2)^2 - 2d\tilde{x}_2d\tilde{x}_3 + (d\tilde{x}_3)^2 \\ (dx_3)^2 = & (d\tilde{x}_3)^2 \end{array}$$

Same geometry, different coordinate system...

$$\begin{bmatrix} -1 & 0 & 0 & 0 \\ 0 & 1 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix} \simeq_{\text{diff}} \begin{bmatrix} -1 & -\cos(\tilde{x}_1) & 0 & 0 \\ -\cos(\tilde{x}_1) & 1 - \cos^2(\tilde{x}_1) & 2\tilde{x}_2 & 0 \\ 0 & 2\tilde{x}_2 & 4\tilde{x}_2^2 + 1 & -1 \\ 0 & 0 & -1 & 2 \end{bmatrix}$$

Consider the change of coordinates  $x_\eta = x_\eta(\tilde{x}^\xi)$  given by:


$$\begin{aligned} x_0 &= \tilde{x}_0 + \sin(\tilde{x}_1) & dx_0 &= d\tilde{x}_0 + \cos(\tilde{x}_1)d\tilde{x}_1 \\ x_1 &= \tilde{x}_1 + \tilde{x}_2^2 & dx_1 &= d\tilde{x}_1 + 2\tilde{x}_2d\tilde{x}_2 \\ x_2 &= \tilde{x}_2 - \tilde{x}_3 & dx_2 &= d\tilde{x}_2 - d\tilde{x}_3 \\ x_3 &= \tilde{x}_3 & dx_3 &= d\tilde{x}_3 \end{aligned} \rightsquigarrow$$

$$\rightsquigarrow \begin{aligned} (dx_0)^2 &= (d\tilde{x}_0)^2 + 2\cos(\tilde{x}_1)d\tilde{x}_0d\tilde{x}_1 + \cos^2(\tilde{x}_1)(d\tilde{x}_1)^2 \\ (dx_1)^2 &= (d\tilde{x}_1)^2 + 4\tilde{x}_2d\tilde{x}_1d\tilde{x}_2 + 4\tilde{x}_2^2(d\tilde{x}_2)^2 \\ (dx_2)^2 &= (d\tilde{x}_2)^2 - 2d\tilde{x}_2d\tilde{x}_3 + (d\tilde{x}_3)^2 \\ (dx_3)^2 &= (d\tilde{x}_3)^2 \end{aligned}$$

Plug to 
$$ds^2 = -(dx_0)^2 + (dx_1)^2 + (dx_2)^2 + (dx_3)^2$$


# Observables

Let  $\mathcal{S}$  be a collection of spacetimes and consider the relation  $\simeq_{\text{diff}}$  on  $\mathcal{S}$ :



# Observables

Let  $\mathcal{S}$  be a collection of spacetimes and consider the relation  $\simeq_{\text{diff}}$  on  $\mathcal{S}$ :




An **observable** is any map  $f: \mathcal{S} \rightarrow R$  that is diffeomorphism invariant:

for all  $g_{\mu\nu}, \tilde{g}_{\rho\sigma} \in \mathcal{S}$  we have  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma} \implies f(g_{\mu\nu}) = f(\tilde{g}_{\rho\sigma})$

# Observables

Let  $\mathcal{S}$  be a collection of spacetimes and consider the relation  $\simeq_{\text{diff}}$  on  $\mathcal{S}$ :



An **observable** is any map  $f: \mathcal{S} \rightarrow R$  that is diffeomorphism invariant:


for all  $g_{\mu\nu}, \tilde{g}_{\rho\sigma} \in \mathcal{S}$  we have  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma} \implies f(g_{\mu\nu}) = f(\tilde{g}_{\rho\sigma})$

Usually  $R = \mathbb{R}$ . For us,  $R$  can be any **Polish space**, such as  $R = \mathbb{R}^{\mathbb{N}}$ .



# Observables

Let  $\mathcal{S}$  be a collection of spacetimes and consider the relation  $\simeq_{\text{diff}}$  on  $\mathcal{S}$ :



An **observable** is any map  $f: \mathcal{S} \rightarrow R$  that is diffeomorphism invariant:

for all  $g_{\mu\nu}, \tilde{g}_{\rho\sigma} \in \mathcal{S}$  we have  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma} \implies f(g_{\mu\nu}) = f(\tilde{g}_{\rho\sigma})$

Usually  $R = \mathbb{R}$ . For us,  $R$  can be any **Polish space**, such as  $R = \mathbb{R}^{\mathbb{N}}$ .

## Canonical Quantization Process

Step 1: Find a **complete** set of observables for  $\mathcal{S}$ .

Step 2: Promote them to an **algebra of operators** on a Hilbert space  $\mathcal{H}$ .

# The problem of observables

*“We define observables as functions (or functionals) of field variables that are invariant with respect to coordinate transformations.”*

(1958) P.G. Bergmann, and A.I. Janis

*“A program aiming at the identification and systematic exploitation of the observables has been under way for many years, but its execution is **hampered by profound technical difficulties, which have not yet been overcome completely.**”*

(1965) P.G. Bergmann,

# The problem of observables

*“We define observables as functions (or functionals) of field variables that are invariant with respect to coordinate transformations.”*

(1958) P.G. Bergmann, and A.I. Janis

*“A program aiming at the identification and systematic exploitation of the observables has been under way for many years, but its execution is **hampered by profound technical difficulties, which have not yet been overcome completely.**”*

(1965) P.G. Bergmann,

*“...presently we can give a formal characterization of observables in general relativity, but we are actually **not able to explicitly construct many examples** of quantities that satisfy it.”*

(2001) L. Smolin

# The problem of observables

*“We define observables as functions (or functionals) of field variables that are invariant with respect to coordinate transformations.”*

(1958) P.G. Bergmann, and A.I. Janis

*“A program aiming at the identification and systematic exploitation of the observables has been under way for many years, but its execution is **hampered by profound technical difficulties, which have not yet been overcome completely.**”*

(1965) P.G. Bergmann,

*“...presently we can give a formal characterization of observables in general relativity, but we are actually **not able to explicitly construct many examples** of quantities that satisfy it.”*

(2001) L. Smolin

*“Observables for full general relativity (without special asymptotic symmetries or matter content) **almost certainly do not exist.**”*


(2015) B. Dittrich, P. A. Höhn, T.A. Koslowski, and M.I. Nelson,

# Examples of Observables

- **Komar mass** for static spacetimes

$$g_{\mu\nu} \mapsto \int_M (2T_{\mu\nu} - Tg_{\mu\nu})u^\mu\xi^\nu dM$$

It is a complete observable for all Schwarzschild solutions



- **ADM Observables** for asymptotically flat spacetimes
- **Coordinate-like Observables** for spacetimes filled with “generic dust”

# Incompleteness of Observables

Theorem (P., Sparling, Christodoulou)

*Complete observables are **not** “analytically definable”*

...in the same way that  $\sqrt[3]{2}$  **cannot** be constructed by “straightedge-and-compass”

# Incompleteness of Observables

Theorem (P., Sparling, Christodoulou)

*Assume that  $\mathcal{S} \supseteq \mathcal{S}_\emptyset$  contains the collection of all **vacuum solutions**  $\mathcal{S}_\emptyset$ .  
Then there is no observable  $f: \mathcal{S} \rightarrow \mathbb{R}$  that is both **Borel** and **complete**.*

# Incompleteness of Observables

Theorem (P., Sparling, Christodoulou)

*Assume that  $\mathcal{S} \supseteq \mathcal{S}_\emptyset$  contains the collection of all **vacuum solutions**  $\mathcal{S}_\emptyset$ . Then there is no observable  $f: \mathcal{S} \rightarrow \mathbb{R}$  that is both **Borel** and **complete**.*

- A **vacuum solution** is any spacetime  $g_{\mu\nu}$  which satisfies:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 0$$



# Incompleteness of Observables

Theorem (P., Sparling, Christodoulou)

Assume that  $\mathcal{S} \supseteq \mathcal{S}_\emptyset$  contains the collection of all **vacuum solutions**  $\mathcal{S}_\emptyset$ .  
Then there is no observable  $f: \mathcal{S} \rightarrow \mathbb{R}$  that is both **Borel** and **complete**.

- A **vacuum solution** is any spacetime  $g_{\mu\nu}$  which satisfies:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 0$$

- $f: \mathcal{S} \rightarrow \mathbb{R}$  is **complete** if  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma} \iff f(g_{\mu\nu}) = f(\tilde{g}_{\rho\sigma})$ .

# Incompleteness of Observables

Theorem (P., Sparling, Christodoulou)

Assume that  $\mathcal{S} \supseteq \mathcal{S}_\emptyset$  contains the collection of all **vacuum solutions**  $\mathcal{S}_\emptyset$ .  
Then there is no observable  $f: \mathcal{S} \rightarrow \mathbb{R}$  that is both **Borel** and **complete**.

- A **vacuum solution** is any spacetime  $g_{\mu\nu}$  which satisfies:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 0$$

- $f: \mathcal{S} \rightarrow \mathbb{R}$  is **complete** if  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma} \iff f(g_{\mu\nu}) = f(\tilde{g}_{\rho\sigma})$ .
- $f: \mathcal{S} \rightarrow \mathbb{R}$  is **Borel** if it is Borel as a map from  $\mathcal{S} \subseteq C^\infty(\mathbb{R}^4, \mathbb{R}^{4 \times 4})$  endowed with the  $C^\infty$ -compact-open topology to the Polish space  $\mathbb{R}$ .

# Incompleteness of Observables

Theorem (P., Sparling, Christodoulou)

Assume that  $\mathcal{S} \supseteq \mathcal{S}_\emptyset$  contains the collection of all **vacuum solutions**  $\mathcal{S}_\emptyset$ .  
Then there is no observable  $f: \mathcal{S} \rightarrow \mathbb{R}$  that is both **Borel** and **complete**.

- A **vacuum solution** is any spacetime  $g_{\mu\nu}$  which satisfies:

$$R_{\mu\nu} - \frac{1}{2}g_{\mu\nu}R = 0$$

- $f: \mathcal{S} \rightarrow \mathbb{R}$  is **complete** if  $g_{\mu\nu} \simeq_{\text{diff}} \tilde{g}_{\rho\sigma} \iff f(g_{\mu\nu}) = f(\tilde{g}_{\rho\sigma})$ .
- $f: \mathcal{S} \rightarrow \mathbb{R}$  is **Borel** if it is Borel as a map from  $\mathcal{S} \subseteq C^\infty(\mathbb{R}^4, \mathbb{R}^{4 \times 4})$  endowed with the  $C^\infty$ -compact-open topology to the Polish space  $\mathbb{R}$ .

Theorem (P., Sparling, Christodoulou)

“ZF+DC+no **complete** observables for  $\mathcal{S} \supseteq \mathcal{S}_\emptyset$  exist” is consistent.

# Table of Contents

1 Spacetimes

2 Observables

3 **The Proof**

4 Future Directions

Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .

Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .

Theorem (Folklore)

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .

**Theorem (Folklore)**

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.

Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .


Theorem (Folklore)

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.
- Assume  $f$  exists. Find comeager  $C \subseteq \{0, 1\}^{\mathbb{Z}}$  so that  $f(C) = \{x\}$





Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .


**Theorem (Folklore)**

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.
- Assume  $f$  exists and get comeager  $C \subseteq \{0, 1\}^{\mathbb{Z}}$  so that  $f(C) = \{x\}$



Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .


**Theorem (Folklore)**

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.
- Assume  $f$  exists and get comeager  $C \subseteq \{0, 1\}^{\mathbb{Z}}$  so that  $f(C) = \{x\}$



Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .


Theorem (Folklore)

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.
- Assume  $f$  exists and get comeager  $C \subseteq \{0, 1\}^{\mathbb{Z}}$  so that  $f(C) = \{x\}$



Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .


**Theorem (Folklore)**

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.
- Assume  $f$  exists and get comeager  $C \subseteq \{0, 1\}^{\mathbb{Z}}$  so that  $f(C) = \{x\}$



Consider the equivalence relation  $\simeq_{\mathbb{Z}}$  on the space  $\{0, 1\}^{\mathbb{Z}}$  where

$$\alpha \simeq_{\mathbb{Z}} \beta \iff \exists k \in \mathbb{Z} \forall n \in \mathbb{Z} \alpha(n+k) = \beta(n)$$

i.e., the **orbit equivalence relation** of the *Bernoulli shift*  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$ .


**Theorem (Folklore)**

There is **no Borel** map  $f: \{0, 1\}^{\mathbb{Z}} \rightarrow R$ , taking values in Polish  $R$ , with

$$\alpha \simeq_{\mathbb{Z}} \beta \iff f(\alpha) = f(\beta), \quad \text{for all } \alpha, \beta \in \{0, 1\}^{\mathbb{Z}}$$

**Proof Sketch.**

- Notice that  $\mathbb{Z} \curvearrowright \{0, 1\}^{\mathbb{Z}}$  has a dense orbit. This implies the “0–1 law”: if  $B \subseteq \{0, 1\}^{\mathbb{Z}}$  is  $\mathbb{Z}$ -invariant and Borel, then one of  $B, B^c$  is comeager.
- Assume  $f$  exists and get comeager  $C \subseteq \{0, 1\}^{\mathbb{Z}}$  so that  $f(C) = \{x\}$



- Since  $\mathbb{Z}$  is countable, there exist  $\alpha \not\simeq_{\mathbb{Z}} \beta$  in  $C$ . But  $f(\alpha) = x = f(\beta)$

# General Strategy

Let  $\mathcal{S}$  be a collection of spacetimes.

In order to prove that:

“there is no observable  $f: \mathcal{S} \rightarrow \mathcal{R}$  that is both **Borel** & **complete**”

it suffices to prove that:

there exists a **Borel reduction** from  $(\{0, 1\}^{\mathbb{Z}}, \simeq_{\mathbb{Z}})$  to  $(\mathcal{S}, \simeq_{\text{diff}})$ ,  
i.e., a Borel map  $r: \{0, 1\}^{\mathbb{Z}} \rightarrow \mathcal{S}$  with  $\alpha \simeq_{\mathbb{Z}} \beta \iff r(\alpha) \simeq_{\text{diff}} r(\beta)$

# General Strategy

Let  $\mathcal{S}$  be a collection of spacetimes.

In order to prove that:

“there is no observable  $f: \mathcal{S} \rightarrow R$  that is both **Borel** & **complete**”

it suffices to prove that:

there exists a **Borel reduction** from  $(\{0, 1\}^{\mathbb{Z}}, \simeq_{\mathbb{Z}})$  to  $(\mathcal{S}, \simeq_{\text{diff}})$ ,  
i.e., a Borel map  $r: \{0, 1\}^{\mathbb{Z}} \rightarrow \mathcal{S}$  with  $\alpha \simeq_{\mathbb{Z}} \beta \iff r(\alpha) \simeq_{\text{diff}} r(\beta)$

## Definition

$\mathcal{S}$  is **rich** if there exists a Borel reduction from  $(\{0, 1\}^{\mathbb{Z}}, \simeq_{\mathbb{Z}})$  to  $(\mathcal{S}, \simeq_{\text{diff}})$

## Examples of Rich Families: part I

Theorem (Christodoulou, Sparling, P.)

*For every  $n \geq 2$ , the family of all spacetimes on  $\mathbb{R}^n$  is rich.*



## Examples of Rich Families: part I

Theorem (Christodoulou, Sparling, P.)

*For every  $n \geq 2$ , the family of all spacetimes on  $\mathbb{R}^n$  is rich.*

**Proof Idea.**

Use the Cosmological Friedmann–Lemaître–Robertson–Walker metrics:

$$g_{\mu\nu} := -dt^2 + W(t)(dx^2 + dy^2 + dz^2)$$

# Examples of Rich Families: part I


Theorem (Christodoulou, Sparling, P.)

For every  $n \geq 2$ , the family of all spacetimes on  $\mathbb{R}^n$  is rich.


## Proof Idea.

Use the Cosmological Friedmann–Lemaître–Robertson–Walker metrics:

$$g_{\mu\nu} := -dt^2 + W(t)(dx^2 + dy^2 + dz^2)$$



Source: Wikipedia



Source: Samuel Velasco/Quanta Magazine

## Examples of Rich Families: part II

Theorem (Christodoulou, Sparling, P.)

*The family  $\mathcal{S}_0$  of all **vacuum solutions** on  $\mathbb{R}^4$  is rich.*

*“The problem already lies in the local degrees of freedom of the background theory in 4D.”*

## Examples of Rich Families: part II

Theorem (Christodoulou, Sparling, P.)

*The family  $\mathcal{S}_0$  of all **vacuum solutions** on  $\mathbb{R}^4$  is rich.*

*“The problem already lies in the local degrees of freedom of the background theory in 4D.”*


**Remark.** There is a unique vacuum solution on  $\mathbb{R}^3$ !

# Proof: Plane Waves

Consider the variables  $u, v, x, y$ .

$$g_{\mu\nu}^H : (u, v, x, y) \mapsto \begin{bmatrix} H(u, x, y) & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$


Is a **vacuum solution** whenever  $H_{xx} + H_{yy} = 0$ .



Penrose: “A Remarkable Property of Plane Waves in General Relativity”

# The reduction

For every  $\alpha \in \{0, 1\}^{\mathbb{Z}}$  we define a “smooth version”  $w^\alpha: \mathbb{R} \rightarrow \mathbb{R}$  of  $\alpha$ :




This defines a map  $r: \{0, 1\}^{\mathbb{Z}} \rightarrow \mathcal{S}_\emptyset$  which maps  $\alpha$  to

$$r(\alpha) := g_{\mu\nu}^\alpha \quad \text{given by} \quad (u, v, x, y) \mapsto \begin{bmatrix} w^\alpha(u)xy & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

# The reduction

For every  $\alpha \in \{0, 1\}^{\mathbb{Z}}$  we define a “smooth version”  $w^\alpha: \mathbb{R} \rightarrow \mathbb{R}$  of  $\alpha$ :




This defines a map  $r: \{0, 1\}^{\mathbb{Z}} \rightarrow \mathcal{S}_\emptyset$  which maps  $\alpha$  to

$$r(\alpha) := g_{\mu\nu}^\alpha \quad \text{given by} \quad (u, v, x, y) \mapsto \begin{bmatrix} w^\alpha(u)xy & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Showing that  $\alpha E_{\mathbb{Z}} \beta \Rightarrow r(\alpha) \simeq_{\text{diff}} r(\beta)$  is easy.

# The reduction

For every  $\alpha \in \{0, 1\}^{\mathbb{Z}}$  we define a “smooth version”  $w^\alpha: \mathbb{R} \rightarrow \mathbb{R}$  of  $\alpha$ :



This defines a map  $r: \{0, 1\}^{\mathbb{Z}} \rightarrow \mathcal{S}_\emptyset$  which maps  $\alpha$  to

$$r(\alpha) := g_{\mu\nu}^\alpha \quad \text{given by} \quad (u, v, x, y) \mapsto \begin{bmatrix} w^\alpha(u)xy & 1 & 0 & 0 \\ 1 & 0 & 0 & 0 \\ 0 & 0 & 1 & 0 \\ 0 & 0 & 0 & 1 \end{bmatrix}$$

- Showing that  $\alpha E_{\mathbb{Z}} \beta \Rightarrow r(\alpha) \simeq_{\text{diff}} r(\beta)$  is easy.
- Showing that  $\alpha E_{\mathbb{Z}} \beta \Leftarrow r(\alpha) \simeq_{\text{diff}} r(\beta)$  is **hard**.



# The difficult direction

**Assume that:**

$$g := (\tilde{w}(\tilde{u})\tilde{x}\tilde{y})d\tilde{u}^2 + 2d\tilde{u}d\tilde{v} + d\tilde{x}^2 + d\tilde{y}^2$$

$$\tilde{g} := (w(u)xy)du^2 + 2dudv + dx^2 + dy^2$$

are diffeomorphic under the smooth change of coordinates  $\varphi$  specified by

$$\tilde{u} = \tilde{u}(u, v, x, y)$$

$$\tilde{v} = \tilde{v}(u, v, x, y)$$

$$\tilde{x} = \tilde{x}(u, v, x, y)$$

$$\tilde{y} = \tilde{y}(u, v, x, y)$$

**Goal:**

To show that  $w(u)$  is a  $\mathbb{Z}$ -shift of  $\tilde{w}(\tilde{u})$ .

# The difficult direction

**Assume that:**

$$g := (\tilde{w}(\tilde{u})\tilde{x}\tilde{y})d\tilde{u}^2 + 2d\tilde{u}d\tilde{v} + d\tilde{x}^2 + d\tilde{y}^2$$

$$\tilde{g} := (w(u)xy)du^2 + 2dudv + dx^2 + dy^2$$

are diffeomorphic under the smooth change of coordinates  $\varphi$  specified by

$$\tilde{u} = \tilde{u}(u, v, x, y)$$

$$\tilde{v} = \tilde{v}(u, v, x, y)$$

$$\tilde{x} = \tilde{x}(u, v, x, y)$$

$$\tilde{y} = \tilde{y}(u, v, x, y)$$

**Goal:**

To show that  $w(u)$  is a  $\mathbb{Z}$ -shift of  $\tilde{w}(\tilde{u})$ .

**Naive approach:** use the definition

$$g_{\mu\nu} = \frac{\partial \tilde{x}^\rho}{\partial x^\mu} \frac{\partial \tilde{x}^\sigma}{\partial x^\nu} \tilde{g}_{\rho\sigma}$$

# Dead end

The relation

$$g_{\mu\nu} = \frac{\partial \tilde{x}^\rho}{\partial x^\mu} \frac{\partial \tilde{x}^\sigma}{\partial x^\nu} \tilde{g}_{\rho\sigma}$$

gives the following equations:

$$\begin{aligned} H(u, x, y) &= \tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_u + 2\tilde{u}_u\tilde{v}_u + \tilde{x}_u^2 + \tilde{y}_u^2 \\ 0 &= \tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_v + 2\tilde{u}_v\tilde{v}_v + \tilde{x}_v^2 + \tilde{y}_v^2 \\ 1 &= \tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_x + 2\tilde{u}_x\tilde{v}_x + \tilde{x}_x^2 + \tilde{y}_x^2 \\ 1 &= \tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_y + 2\tilde{u}_y\tilde{v}_y + \tilde{x}_y^2 + \tilde{y}_y^2 \\ 1 &= 2\tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_u\tilde{u}_v + 2(\tilde{u}_u\tilde{v}_v + \tilde{u}_v\tilde{v}_u) + 2\tilde{x}_u\tilde{x}_v + 2\tilde{y}_u\tilde{y}_v \\ 0 &= 2\tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_x\tilde{u}_y + 2(\tilde{u}_x\tilde{v}_y + \tilde{u}_y\tilde{v}_x) + 2\tilde{x}_x\tilde{x}_y + 2\tilde{y}_x\tilde{y}_y \\ 0 &= 2\tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_u\tilde{u}_x + 2(\tilde{u}_u\tilde{v}_x + \tilde{u}_x\tilde{v}_u) + 2\tilde{x}_u\tilde{x}_x + 2\tilde{y}_u\tilde{y}_x \\ 0 &= 2\tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_u\tilde{u}_y + 2(\tilde{u}_u\tilde{v}_y + \tilde{u}_y\tilde{v}_u) + 2\tilde{x}_u\tilde{x}_y + 2\tilde{y}_u\tilde{y}_y \\ 0 &= 2\tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_v\tilde{u}_x + 2(\tilde{u}_v\tilde{v}_x + \tilde{u}_x\tilde{v}_v) + 2\tilde{x}_v\tilde{x}_x + 2\tilde{y}_v\tilde{y}_x \\ 0 &= 2\tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})\tilde{u}_v\tilde{u}_y + 2(\tilde{u}_v\tilde{v}_y + \tilde{u}_y\tilde{v}_v) + 2\tilde{x}_v\tilde{x}_y + 2\tilde{y}_v\tilde{y}_y \end{aligned}$$

Good Luck!

Instead: analyze the Killing vector fields!

By analyzing the Lie algebra of Killing fields: every diffeo  $\varphi$  between

$$g = \tilde{H}(\tilde{u}, \tilde{x}, \tilde{y})d\tilde{u}^2 + 2d\tilde{u}d\tilde{v} + d\tilde{x}^2 + d\tilde{y}^2$$

$$\tilde{g} = H(u, x, y)du^2 + 2dudv + dx^2 + dy^2$$

has to be of the following form, for some  $a, b, c$  and  $f(u), g(u), h(u)$ :

$$\begin{aligned}\tilde{u} &= (u + a)/c \\ \tilde{x} &= x \cos(b) + y \sin(b) + g(u) \\ \tilde{y} &= -x \sin(b) + y \cos(b) + h(u) \\ \tilde{v} &= c[v - x(\cos(b)g'(u) - \sin(b)h'(u)) \\ &\quad - y(\sin(b)g'(u) - \cos(b)h'(u)) - f(u)]\end{aligned}$$

Jordan, Ehlers, Kundt (based on work of Robinson)

# Table of Contents

1 Spacetimes


2 Observables

3 The Proof

4 Future Directions

## Canonical Quantization

Step 1: Find **complete** set of observables.




Step 2: promote them to an **algebra of operators** on a Hilbert space  $\mathcal{H}$ .

## Canonical Quantization

~~Step 1: Find **complete** set of observables.~~

Cannot be done in a constructive fashion if  $\mathcal{S} \supseteq \{\text{vacuum solutions}\}$




Step 2: promote them to an **algebra of operators** on a Hilbert space  $\mathcal{H}$ .

## Canonical Quantization


~~Step 1: Find **complete** set of observables.~~

Cannot be done in a constructive fashion if  $\mathcal{S} \supseteq \{\text{vacuum solutions}\}$



Step 2: promote them to an **algebra of operators** on a Hilbert space  $\mathcal{H}$ .


## Equivariant forms of Quantization



$G$ -observables where  $G$  has **nice representation theoretic properties**.




# The Borel reduction hierarchy




A **classification problem**  $(X, E)$  is an equivalence relation  $E$  on Polish  $X$

$(X, E) \leq (\tilde{X}, \tilde{E})$  iff  $(X, E)$  **Borel reduces** to  $(Y, F)$   
iff there exists Borel  $r: X \rightarrow Y$  so that  $xEx' \iff r(x)Fr(x')$

**Program.** Place  $(\mathcal{S}, \simeq_{\text{diff}})$  in the **Borel reduction hierarchy**



**Program.** Place  $(\mathcal{S}, \simeq_{\text{diff}})$  in the **Borel reduction hierarchy**



Thank you!